

SA's Leading Past Year

Exam Paper Portal



You have Downloaded, yet Another Great Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexampapers.co.za



**SA EXAM
PAPERS**
SA EXAM
PAPERS



basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

**NATIONAL
SENIOR CERTIFICATE/
NASIONALE
SENIOR SERTIFIKAAT**

GRADE 12/*GRAAD 12*

MATHEMATICS P2/*WISKUNDE V2*

NOVEMBER 2023

MARKING GUIDELINES/*NASIENRIGLYNE*

MARKS/PUNTE: 150

**These marking guidelines consist of 23 pages./
*Hierdie nasienriglyne bestaan uit 23 bladsye.***



NOTE:

- If a candidate answers a question TWICE, only mark the FIRST attempt.
- If a candidate has crossed out an attempt of a question and not redone the question, mark the crossed out version.
- Consistent accuracy applies in ALL aspects of the Marking Guidelines. Stop marking at the second calculation error.
- Assuming answers/values in order to solve a problem is NOT acceptable.

NOTA:

- *As 'n kandidaat 'n vraag TWEE KEER beantwoord, merk slegs die EERSTE poging.*
- *As 'n kandidaat 'n antwoord van 'n vraag doodtrek en nie oordoen nie, merk die doodgetrekte poging.*
- *Volgehoue akkuraatheid word in ALLE aspekte van die Nasienriglyne toegepas. Hou op nasien by die tweede berekeningsfout.*
- *Aanvaar van antwoorde/waardes om 'n probleem op te los, word NIE toegelaat nie.*

| GEOMETRY | |
|-----------------|---|
| S | A mark for a correct statement (A statement mark is independent of a reason) |
| | 'n Punt vir 'n korrekte bewering ('n Punt vir 'n bewering is onafhanklik van die rede) |
| R | A mark for the correct reason (A reason mark may only be awarded if the statement is correct) |
| | 'n Punt vir 'n korrekte rede ('n Punt word slegs vir die rede toegeken as die bewering korrek is) |
| S/R | Award a mark if statement AND reason are both correct |
| | Ken 'n punt toe as die bewering EN rede beide korrek is |



QUESTION/VRAAG 1

| | | |
|-----|---|---|
| 1.1 | $a = -23,846\dots$ $b = 0,227\dots$ $\hat{y} = -23,85 + 0,23x$ | ✓ $a = -23,846\dots$ ✓ $b = 0,227\dots$ ✓ equation (3) |
| 1.2 | $\hat{y} = -23,85 + 0,23(550)$ $y = 102,65$ OR $y = 101,02$ | ✓ substitution of 550 ✓ answer (2) ✓✓ $y = 101,02$ (calculator) (2) |
| 1.3 | $r = 0,98$ | ✓ $r = 0,98$ (1) |
| 1.4 | Very strong positive correlation | ✓ strong positive (1) |

| | | | | | | | |
|----|-----|-----|-----|-----|-----|-----|-----|
| 50 | 100 | 130 | 150 | 180 | 190 | 200 | 200 |
|----|-----|-----|-----|-----|-----|-----|-----|

| | | |
|-------|---|--|
| 1.5.1 | $\bar{x} = \frac{1200}{8}$ $\bar{x} = 150$ OR $\bar{x} = 150$ | ✓ 1200 ✓ answer (2) ✓✓ $\bar{x} = 150$ (2) |
| 1.5.2 | $\sigma = 50,50$ | ✓ $\sigma = 50,50$ (1) |
| 1.5.3 | $\bar{x} - \sigma$ $= 150 - 50,50$ $= 99,50$ \therefore 1 stop | ✓ calculation of $\bar{x} - \sigma$ ✓ answer (2) |
| | | [12] |




QUESTION/VRAAG 2

| 2.1 | <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>Number of glasses of water per day</th> <th>Number of staff members</th> <th>Cumulative frequency</th> </tr> </thead> <tbody> <tr> <td>$0 \leq x < 2$</td> <td>5</td> <td>5</td> </tr> <tr> <td>$2 \leq x < 4$</td> <td>15</td> <td>20</td> </tr> <tr> <td>$4 \leq x < 6$</td> <td>13</td> <td>33</td> </tr> <tr> <td>$6 \leq x < 8$</td> <td>5</td> <td>38</td> </tr> <tr> <td>$8 \leq x < 10$</td> <td>2</td> <td>40</td> </tr> </tbody> </table> | Number of glasses of water per day | Number of staff members | Cumulative frequency | $0 \leq x < 2$ | 5 | 5 | $2 \leq x < 4$ | 15 | 20 | $4 \leq x < 6$ | 13 | 33 | $6 \leq x < 8$ | 5 | 38 | $8 \leq x < 10$ | 2 | 40 | <p>✓ 5; 20</p> <p>✓ 40</p> <p style="text-align: right;">(2)</p> |
|------------------------------------|--|--|-------------------------|----------------------|----------------|---|---|----------------|----|----|----------------|----|----|----------------|---|----|-----------------|---|----|--|
| Number of glasses of water per day | Number of staff members | Cumulative frequency | | | | | | | | | | | | | | | | | | |
| $0 \leq x < 2$ | 5 | 5 | | | | | | | | | | | | | | | | | | |
| $2 \leq x < 4$ | 15 | 20 | | | | | | | | | | | | | | | | | | |
| $4 \leq x < 6$ | 13 | 33 | | | | | | | | | | | | | | | | | | |
| $6 \leq x < 8$ | 5 | 38 | | | | | | | | | | | | | | | | | | |
| $8 \leq x < 10$ | 2 | 40 | | | | | | | | | | | | | | | | | | |
| 2.2 | 40 staff members | <p>✓ answer</p> <p style="text-align: right;">(1)</p> | | | | | | | | | | | | | | | | | | |
| 2.3 | 33 staff members | <p>✓ answer</p> <p style="text-align: right;">(1)</p> | | | | | | | | | | | | | | | | | | |
| 2.4 | $\bar{x} = \frac{\left(1 \times \left(5 + \frac{k}{2}\right)\right) + (3 \times 15) + \left(5 \times \left(13 + \frac{k}{2}\right)\right) + (7 \times 5) + (9 \times 2)}{40 + k} = 4$ $5 + \frac{k}{2} + 45 + 65 + \frac{5k}{2} + 35 + 18 = 160 + 4k$ $3k + 168 = 160 + 4k$ $k = 8$ <p>OR</p> $\bar{x} = \frac{(1 \times 5) + (15 \times 3) + (13 \times 5) + (5 \times 7) + (2 \times 9)}{40}$ $= 4,2$ $\bar{x}_{\text{old}} - \bar{x}_{\text{current}} = 4,2 - 4$ $= 0,2$ $\therefore 0,2 \times 40$ $= 8 \text{ teachers}$ | <p>✓ answer from Q2.2 + k</p> <p>✓ $\left(1 \times \left(5 + \frac{k}{2}\right)\right)$</p> <p>✓ $\left(5 \times \left(13 + \frac{k}{2}\right)\right)$</p> <p>✓ answer</p> <p style="text-align: right;">(4)</p> <p>✓ 4,2</p> <p>✓ $\bar{x}_{\text{old}} - 4$</p> <p>✓ difference</p> <p>✓ answer</p> <p style="text-align: right;">(4)</p> | | | | | | | | | | | | | | | | | | |
| | | [8] | | | | | | | | | | | | | | | | | | |



QUESTION/VRAAG 3



| | | |
|-----|---|---|
| 3.1 | $SL = \sqrt{(x_s - x_L)^2 + (y_s - y_L)^2}$ $SL = \sqrt{(4 - (-4))^2 + (5 - 1)^2}$ $SL = \sqrt{80} = 4\sqrt{5} = 8,94 \text{ units}$ | ✓ substitution of S and L into correct formula ✓ answer (2) |
| 3.2 | $m_{SN} = \frac{5 - (-3)}{4 - (-2)}$ $m_{SN} = \frac{4}{3}$ | ✓ substitution of S and N into correct formula ✓ answer (2) |
| 3.3 | $m = \tan \theta = \frac{4}{3}$ $\theta = 53,13^\circ$ | ✓ $\tan \theta = m_{SN}$ ✓ answer (2) |
| 3.4 | $m_{LN} = \frac{1 - (-3)}{-4 - (-2)}$ $m_{LN} = -2$ $\hat{L}\hat{K}\hat{O} = 116,565\dots^\circ$ $\hat{L}\hat{N}\hat{S} = 116,565\dots^\circ - 53,13^\circ$ $\hat{L}\hat{N}\hat{S} = 63,44^\circ$ | ✓ $m_{LN} = -2$ ✓ size of $\hat{L}\hat{K}\hat{O}$ ✓ answer (3) |




| | | |
|-----|---|--|
| | <p>OR</p> <p>SN = 10 units $\sin \hat{LNS} = \frac{4\sqrt{5}}{10}$ $\hat{LNS} = 63,44^\circ$</p> <p>OR</p> <p>LN = $2\sqrt{5}$ units $\tan \hat{LNS} = \frac{4\sqrt{5}}{2\sqrt{5}}$ $\hat{LNS} = 63,44^\circ$</p> <p>OR</p> <p>SN = 10 units LN = $2\sqrt{5}$ units $\cos \hat{LNS} = \frac{2\sqrt{5}}{10}$ $\hat{LNS} = 63,44^\circ$</p> | <p>✓ SN = 10 units</p> <p>✓ correct trig ratio</p> <p>✓ answer (3)</p> <p>✓ LN = $2\sqrt{5}$ units</p> <p>✓ correct trig ratio</p> <p>✓ answer (3)</p> <p>✓ SN = 10 units and LN = $2\sqrt{5}$ units</p> <p>✓ correct trig ratio</p> <p>✓ answer (3)</p> |
| 3.5 | <p>$m = \frac{4}{3}$</p> <p>$1 = \frac{4}{3}(-4) + c$ OR $y - 1 = \frac{4}{3}(x - (-4))$</p> <p>$c = \frac{19}{3}$ $y - 1 = \frac{4}{3}x + \frac{16}{3}$</p> <p>$y = \frac{4}{3}x + \frac{19}{3}$ $y = \frac{4}{3}x + \frac{19}{3}$</p> | <p>✓ m_{SN}</p> <p>✓ substitution of m_{SN} & L</p> <p>✓ equation (3)</p> |
| 3.6 | <p>SL = $4\sqrt{5}$</p> <p>LN = $\sqrt{(-4 - (-2))^2 + (1 - (-3))^2}$</p> <p>LN = $\sqrt{20} = 2\sqrt{5}$</p> <p>Area $\Delta LSN = \frac{1}{2}(4\sqrt{5})(2\sqrt{5})$ = 20 units²</p> <p>OR</p> | <p>✓ LN = $\sqrt{20} = 2\sqrt{5}$</p> <p>✓ substitution into formula</p> <p>✓ answer (3)</p> |



| | | |
|------------|---|---|
| | <p>SN = 10 units</p> $LN = \sqrt{(-4 - (-2))^2 + (1 - (-3))^2}$ $LN = \sqrt{20} = 2\sqrt{5}$ $\text{Area } \triangle LSN = \frac{1}{2}(10)(2\sqrt{5})\sin 63,44^\circ$ $= 20 \text{ units}^2$ | <p>✓ $LN = \sqrt{20} = 2\sqrt{5}$</p> <p>✓ substitution into formula</p> <p>✓ answer</p> <p>(3)</p> |
| <p>3.7</p> | <p>$\hat{L} = 90^\circ$ SN is a diameter of circle S, L, N [chord subtends 90° OR converse \angle in semi-circle]</p> <p>Centre of circle = P $\left(\frac{4 + (-2)}{2}; \frac{5 + (-3)}{2}\right)$ $= P(1; 1)$</p> <p>OR Let the coordinates of P be $(a; b)$. Then, PL = PN: $(-4 - a)^2 + (1 - b)^2 = (-2 - a)^2 + (-3 - b)^2$ $a - 2b = -1$equation 1 If PS = PN, then: $4a + 2b = 6$ equation 2 Solving simultaneously yields: $a = 1$ and $b = 1$ and P(1; 1)</p> <p>OR If PL = PN, then: $a - 2b = -1$equation 1 If PS = PL, then: $2a + b = 3$equation 2 Solving simultaneously yields: $a = 1$ and $b = 1$ and P(1; 1)</p> | <p>✓ SN is a diameter of circle S, L, N</p> <p>✓ x-value ✓ y-value</p> <p>(3)</p> <p>✓ 2 correct linear equations ✓ x-value ✓ y-value</p> <p>(3)</p> <p>✓ 2 correct linear equations ✓ x-value ✓ y-value</p> <p>(3)</p> |
| <p>3.8</p> | <p>$\hat{LPN} = \theta = 53,13^\circ$ [alt \angles; LP \parallel x-axis] $\therefore \hat{LPS} = 126,87^\circ$</p> <p>OR</p> <p>$\hat{LNS} = 63,44^\circ$ $\therefore \hat{LPS} = 126,88^\circ$ [\angle at centre = $2 \times \angle$ at circumference]</p> <p>OR</p> <p>$\hat{LSN} = 26,56^\circ$ [sum of \angles in Δ] $\hat{SLP} = 26,56^\circ$ [\angles opp equal radii] $\therefore \hat{LPS} = 126,88^\circ$ [sum of \angles in Δ]</p> <p>OR</p> <p>$(4\sqrt{5})^2 = 5^2 + 5^2 - 2(5)(5)\cos \hat{LPS}$ $\cos \hat{LPS} = -\frac{3}{5}$ $\therefore \hat{LPS} = 126,87^\circ$</p> | <p>✓ \hat{LPN}</p> <p>✓ answer</p> <p>(2)</p> <p>✓ \hat{LNS}</p> <p>✓ answer</p> <p>(2)</p> <p>✓ \hat{LSN}</p> <p>✓ answer</p> <p>(2)</p> <p>✓ correct substitution into cosine formula</p> <p>✓ answer</p> <p>(2)</p> |
| | | <p>[20]</p> |



QUESTION/VRAAG 4

| | | | | |
|-----|--|---|---------------------------------------|-----|
| 4.1 | $D(p; -2)$ $x^2 + y^2 = 20$ $p^2 + (-2)^2 = 20$ $p^2 = 16$ $p = \pm 4$ $p = 4$ | ✓ substitution of point $D(p; -2)$ ✓ $p^2 = 16$ | (2) | |
| 4.2 | $\frac{4 + x_F}{2} = 6$ $x_F = 8$ $F(8; 6)$ OR $x_E - x_D = 6 - 4$ $= 2$ $x_F = 6 + 2 = 8$ $F(8; 6)$ | $\frac{-2 + y_F}{2} = 2$ $y_F = 6$ $y_E - y_D = 2 - (-2)$ $= 4$ $y_F = 2 + 4 = 6$ | ✓ method ✓ x -value ✓ y -value | (3) |
| | | ✓ method ✓ x -value ✓ y -value | (3) | |




| | | |
|-----|--|---|
| 4.3 | $m_{DE} = \frac{-2-2}{4-6}$ $m_{DE} = 2$ $-2 = 2(4) + c$ $c = -10$ $y = 2x - 10$ <p>OR</p> $m_{OD} = -\frac{2}{4} = -\frac{1}{2}$ $\therefore m_{DE} = 2$ $-2 = 2(4) + c$ $c = -10$ $y = 2x - 10$ <p>OR</p> $y - (-2) = 2(x - 4)$ $y + 2 = 2x - 8$ $y = 2x - 10$ | <ul style="list-style-type: none"> ✓ correct substitution ✓ gradient of DE, DF or EF ✓ substitution of point D(4 ; -2) or E(6 ; 2) or F(8 ; 6) ✓ answer (4) ✓ correct gradient of OD ✓ gradient of DE [tan ⊥ radius] ✓ substitution of point D(4 ; -2) or E(6 ; 2) or F(8 ; 6) ✓ answer (4) |
| 4.4 | $m_{DE} = 2$ $\therefore m_{GF} = -\frac{1}{2}$ $\frac{0-6}{t-8} = -\frac{1}{2}$ $-(t-8) = 2(-6)$ $t = 20$ <p>OR</p> $y = 2x - 10$ $0 = 2x - 10$ $x = 5$ $A(5 ; 0)$ <p>In $\triangle AFG$: $FA \perp FG$</p> $FA^2 = (6-0)^2 + (8-5)^2 = 45$ $FG^2 = (t-8)^2 + (0-6)^2$ $= t^2 - 16t + 100$ $GA^2 = (t-5)^2$ $= t^2 - 10t + 25$ $\therefore GA^2 = GF^2 + FA^2$ $t^2 - 10t + 25 = t^2 - 16t + 100 + 45$ $6t = 120$ $t = 20$ | <ul style="list-style-type: none"> ✓ correct gradient of GF ✓ substitution of F ✓ answer (3) ✓ x-intercept of DF ✓ substitution into Pythagoras ✓ answer (3) |

| | | |
|-------------|--|--|
| 4.5 | $F(8;6)$ $G(20;0)$ $(8-20)^2 + (6-0)^2 = r^2$ $r^2 = 180$ $(x-20)^2 + y^2 = 180$ $x^2 + y^2 - 40x + 220 = 0$ | ✓ substitution of F and G ✓ value of r^2 ✓ equation of circle ✓ answer (4) |
| 4.6 | <p>Smaller circle $r = 2\sqrt{5}$ Larger circle $r = 6\sqrt{5}$</p> $G(20;0)$ $k = 20 - (6\sqrt{5} - 2\sqrt{5}) \quad \text{or} \quad k = 20 + (6\sqrt{5} - 2\sqrt{5})$ $= 20 - 4\sqrt{5} \quad \quad \quad = 20 + 4\sqrt{5}$ $= 11,06 \text{ units} \quad \quad \quad = 28,94 \text{ units}$ <p>OR</p> <p>Smaller circle $r = 2\sqrt{5}$</p> $k = 2(2\sqrt{5}) + 20 - 8\sqrt{5} \quad \text{or} \quad k = 2(6\sqrt{5}) + 20 - 8\sqrt{5}$ $= 20 - 4\sqrt{5} \quad \quad \quad = 20 + 4\sqrt{5}$ $= 11,06 \text{ units} \quad \quad \quad = 28,94 \text{ units}$ <p>OR</p> $x^2 + y^2 - 40x + 220 = 0$ $y = 0$ $\therefore x^2 - 40x + 220 = 0$ $\therefore x = 20 + 6\sqrt{5} \quad \text{or} \quad x = 20 - 6\sqrt{5}$ $\therefore k = 20 + 6\sqrt{5} - \sqrt{20} \quad \text{or} \quad k = 20 - 6\sqrt{5} + \sqrt{20}$ $\therefore k = 20 + 4\sqrt{5} \quad \quad \quad \therefore k = 20 - 4\sqrt{5}$ $= 11,06 \text{ units} \quad \quad \quad = 28,94 \text{ units}$ | ✓ $r = 2\sqrt{5}$ ✓ method ✓ answer ✓ answer (4) |
| | | ✓ $r = 2\sqrt{5}$ ✓ method ✓ answer ✓ answer (4) |
| [20] | | |



QUESTION/VRAAG 5

| | | |
|-------|--|--|
| 5.1.1 | $\sin \beta = \frac{1}{3} \quad \beta \in (90^\circ; 270^\circ)$  $x = -\sqrt{8} = -2\sqrt{2}$ $\cos \beta = \frac{-2\sqrt{2}}{3}$ <p>OR</p> $\sin \beta = \frac{1}{3} \quad \beta \in (90^\circ; 270^\circ)$ $\cos^2 \beta = 1 - \sin^2 \beta$ $\cos^2 \beta = 1 - \left(\frac{1}{3}\right)^2$ $\cos^2 \beta = \frac{8}{9}$ $\cos \beta = \frac{-\sqrt{8}}{3} = \frac{-2\sqrt{2}}{3}$ | <p>✓ $x^2 + y^2 = r^2$</p> <p>✓ $x = -2\sqrt{2}$</p> <p>✓ answer (3)</p> <p>✓ square identity</p> <p>✓ $\cos^2 \beta$</p> <p>✓ answer (3)</p> |
| 5.1.2 | $\sin 2\beta = 2 \sin \beta \cos \beta$ $= 2 \left(\frac{1}{3}\right) \left(\frac{-\sqrt{8}}{3}\right)$ $= \frac{-2\sqrt{8}}{9} \quad \text{OR} \quad 2 \left(\frac{-2\sqrt{2}}{9}\right)$ $= \frac{-4\sqrt{2}}{9}$ | <p>✓ double angle</p> <p>✓ substitution</p> <p>✓ answer (3)</p> |
| 5.1.3 | $\cos (450^\circ - \beta)$ $= \cos (90^\circ - \beta)$ $= \sin \beta$ $= \frac{1}{3}$ <p>OR</p> | <p>✓ $\cos (90^\circ - \beta)$</p> <p>✓ co-ratio</p> <p>✓ answer (3)</p> |


| | | | |
|-------|---|--|-----|
| | $\begin{aligned} & \cos(450^\circ - \beta) \\ &= \cos 450^\circ \cos \beta + \sin 450^\circ \sin \beta \\ &= \cos 90^\circ \cos \beta + \sin 90^\circ \sin \beta \\ &= \sin \beta \\ &= \frac{1}{3} \end{aligned}$ | <ul style="list-style-type: none"> ✓ expansion ✓ reduction ✓ answer | (3) |
| 5.2.1 | $\begin{aligned} \text{LHS} &= \frac{\cos^4 x + \sin^2 x \cdot \cos^2 x}{1 + \sin x} \\ &= \frac{\cos^2 x (\cos^2 x + \sin^2 x)}{1 + \sin x} \\ &= \frac{1 - \sin^2 x}{1 + \sin x} \\ &= \frac{(1 - \sin x)(1 + \sin x)}{1 + \sin x} \\ &= 1 - \sin x \\ &= \text{RHS} \end{aligned}$ <p>OR</p> $\begin{aligned} \text{LHS} &= \frac{\cos^4 x + \sin^2 x \cdot \cos^2 x}{1 + \sin x} \\ &= \frac{\cos^4 x + (1 - \cos^2 x) \cos^2 x}{1 + \sin x} \\ &= \frac{\cos^4 x + \cos^2 x - \cos^4 x}{1 + \sin x} \\ &= \frac{1 - \sin^2 x}{1 + \sin x} \\ &= \frac{(1 - \sin x)(1 + \sin x)}{1 + \sin x} \\ &= 1 - \sin x \\ &= \text{RHS} \end{aligned}$ <p>OR</p> $\begin{aligned} \text{RHS} &= 1 - \sin x \\ &= (1 - \sin x) \times \frac{1 + \sin x}{1 + \sin x} \\ &= \frac{1 - \sin^2 x}{1 + \sin x} \\ &= \frac{\cos^2 x}{1 + \sin x} \\ &= \frac{\cos^2 x (\sin^2 x + \cos^2 x)}{1 + \sin x} \\ &= \frac{\cos^4 x + \cos^2 x \cdot \sin^2 x}{1 + \sin x} \\ &= \text{LHS} \end{aligned}$ | <ul style="list-style-type: none"> ✓ factors ✓ $\sin^2 x + \cos^2 x = 1$ ✓ $\cos^2 x = 1 - \sin^2 x$ ✓ factors ✓ $\sin^2 x = 1 - \cos^2 x$ ✓ expansion ✓ $\cos^2 x = 1 - \sin^2 x$ ✓ factors ✓ $\times \frac{1 + \sin x}{1 + \sin x}$ ✓ product ✓ $1 - \sin^2 x = \cos^2 x$ ✓ $1 = \cos^2 x + \sin^2 x$ | (4) |
| | | | (4) |

| | | |
|-------|---|--|
| 5.2.2 | $\sin x + 1 = 0$ $\sin x = -1$ ref. $\angle = 90^\circ$ $x = 270^\circ$ | $\checkmark \sin x + 1 = 0$ $\checkmark x = 270^\circ$ (2) |
| 5.2.3 | $y = \frac{\cos^4 x + \sin^2 x \cdot \cos^2 x}{1 + \sin x}$ $y = 1 - \sin x$ $\therefore \text{Minimum} = 0$ | $\checkmark \checkmark \text{Minimum} = 0$ (2) |
| 5.3.1 | $\sin(A - B)$ $= \cos[90^\circ - (A - B)]$ $= \cos[(90^\circ - A) - (-B)]$ $= \cos(90^\circ - A)\cos(-B) + \sin(90^\circ - A)\sin(-B)$ $= \sin A \cos B + \cos A(-\sin B)$ $= \sin A \cos B - \cos A \sin B$ OR $\sin(A - B)$ $= \cos[90^\circ - (A - B)]$ $= \cos[(90^\circ + B) - A]$ $= \cos(90^\circ + B)\cos A + \sin(90^\circ + B)\sin A$ $= -\sin B \cos A + \cos B \sin A$ $= \sin A \cos B - \cos A \sin B$ | \checkmark co-ratio \checkmark compound angle \checkmark reduction (3) \checkmark co-ratio \checkmark compound angle \checkmark reduction (3) |
| 5.3.2 | $\sin 48^\circ \cos x - \cos 48^\circ \sin x = \cos 2x$ $\sin(48^\circ - x) = \cos 2x$ $\sin(48^\circ - x) = \sin(90^\circ - 2x)$ $48^\circ - x = 90^\circ - 2x + k \cdot 360^\circ$ or $48^\circ - x = 180^\circ - (90^\circ - 2x) + k \cdot 360^\circ$ $x = 42^\circ + k \cdot 360^\circ$ | \checkmark compound angle \checkmark co-ratio \checkmark both equations \checkmark general solution \checkmark general solution; $k \in \mathbb{Z}$ (5) |
| | OR $\sin 48^\circ \cos x - \cos 48^\circ \sin x = \cos 2x$ $\sin(48^\circ - x) = \cos 2x$ $\cos(90^\circ - 48^\circ + x) = \cos 2x$ $\cos(42^\circ + x) = \cos 2x$ $42^\circ + x = 2x + k \cdot 360^\circ$ or $42^\circ + x = 360^\circ - 2x + k \cdot 360^\circ$ $-x = -42^\circ + k \cdot 360^\circ$ $3x = 318^\circ + k \cdot 360^\circ$ $x = 42^\circ - k \cdot 360^\circ$ $x = 106^\circ + k \cdot 120^\circ$; $k \in \mathbb{Z}$ | \checkmark compound angle \checkmark co-ratio \checkmark both equations \checkmark general solution \checkmark general solution; $k \in \mathbb{Z}$ (5) |


| | | |
|-----|---|---|
| 5.4 | $\frac{\sin 3x + \sin x}{\cos 2x + 1}$ $= \frac{\sin(2x + x) + \sin(2x - x)}{\cos 2x + 1}$ $= \frac{\sin 2x \cos x + \cos 2x \sin x + \sin 2x \cos x - \cos 2x \sin x}{2 \cos^2 x - 1 + 1}$ $= \frac{2 \sin 2x \cos x}{2 \cos^2 x}$ $= \frac{2(2 \sin x \cos x) \cos x}{2 \cos^2 x}$ $= \frac{4 \sin x \cos^2 x}{2 \cos^2 x}$ $= 2 \sin x$ <p>OR</p> $\frac{\sin 3x + \sin x}{\cos 2x + 1}$ $= \frac{\sin(2x + x) + \sin x}{2 \cos^2 x - 1 + 1}$ $= \frac{\sin 2x \cos x + \cos 2x \sin x + \sin x}{2 \cos^2 x}$ $= \frac{2 \sin x \cos x \cos x + \cos 2x \sin x + \sin x}{2 \cos^2 x}$ $= \frac{\sin x(2 \cos^2 x + \cos 2x + 1)}{2 \cos^2 x}$ $= \frac{\sin x(2 \cos^2 x + 2 \cos^2 x - 1 + 1)}{2 \cos^2 x}$ $= 2 \sin x$ | <p>✓ $3x = (2x + x)$</p> <p>✓ expansion</p> <p>✓ double angle of $\cos 2x$</p> <p>✓ simplification</p> <p>✓ $\sin 2x = 2 \sin x \cos x$</p> <p>✓ answer (6)</p> <p>✓ $3x = (2x + x)$</p> <p>✓ double angle of $\cos 2x$</p> <p>✓ expansion</p> <p>✓ $\sin 2x = 2 \sin x \cos x$</p> <p>✓ common factor</p> <p>✓ answer (6)</p> |
| | | [31] |




QUESTION/VRAAG 6




| | | |
|-------|--|--|
| 6.1 | Period = 180° | ✓ 180° (1) |
| 6.2 | $y \in \left[-\frac{\sqrt{2}}{2}; 1\right]$ OR $y \in [-0,71; 1]$ OR $-\frac{\sqrt{2}}{2} \leq y \leq 1$ | ✓ $-\frac{\sqrt{2}}{2}$ ✓ $y \in \left[-\frac{\sqrt{2}}{2}; 1\right]$ (2) |
| 6.3.1 | $x \in (45^\circ; 90^\circ)$ OR $45^\circ < x < 90^\circ$ | ✓✓ $x \in (45^\circ; 90^\circ)$ (2) |
| 6.3.2 | $f(x) + 1 \leq 0$ $f(x) \leq -1$ $x \in [105^\circ; 165^\circ]$ OR $105^\circ \leq x \leq 165^\circ$ | ✓✓ $x \in [105^\circ; 165^\circ]$ (2) |
| 6.4 | $p(x) = -2 \sin 2x$ $-2 \sin 2x = -1$ OR $2 \sin 2x = 1$ $k = 15^\circ$ or $k = 75^\circ$ | ✓ reading off $f(x) = 1$ or $-f(x) = -1$ ✓ 15° ✓ 75° (3) |
| 6.5 | $g(x) = -\cos(x + 45^\circ)$ $h(x) = -\cos(x + 90^\circ)$ $h(x) = \sin x$ | ✓ $-\cos(x + 90^\circ)$ ✓ answer (2) |
| | | [12] |



QUESTION/VRAAG 7




| | | | |
|-----|--|---|-----|
| 7.1 | $\text{Area } \triangle STK = \frac{1}{2} p(\text{SK}) \sin \alpha$ $q = \frac{1}{2} p(\text{SK}) \sin \alpha$ $\text{SK} = \frac{q}{\frac{1}{2} p \sin \alpha}$ $= \frac{2q}{p \sin \alpha}$ | <ul style="list-style-type: none"> ✓ substitution into the correct formula ✓ answer | (2) |
| 7.2 | $\hat{RKS} = \beta$ $\frac{RS}{SK} = \tan \beta$ $RS = \frac{2q \tan \beta}{p \sin \alpha}$ <p>OR</p> $\frac{RS}{\sin \beta} = \frac{SK}{\sin(90^\circ - \beta)}$ $RS \cos \beta = SK \sin \beta$ $RS = SK \tan \beta$ $RS = \frac{2q \tan \beta}{p \sin \alpha}$ | <ul style="list-style-type: none"> ✓ $\hat{RKS} = \beta$ ✓ correct trig ratio | (2) |
| 7.3 | $70 = \frac{2(2500) \tan 42^\circ}{80 \sin \alpha}$ $\sin \alpha = \frac{25}{28} \tan 42^\circ \quad \text{OR} \quad \sin \alpha = 0,80\dots$ $\alpha = 53,51^\circ$ | <ul style="list-style-type: none"> ✓ correct substitution of values into RS ✓ value of $\sin \alpha$ ✓ answer | (3) |
| | | | [7] |



QUESTION/VRAAG 8


8.1



| | | |
|------------|--|--|
| <p>8.1</p> | <p>Construction: Draw KO produced</p> $\hat{O}_1 = \hat{K}_1 + \hat{T} \quad [\text{ext } \angle \text{ of } \Delta]$ <p>But $\hat{K}_1 = \hat{T}$ [\angles opp equal sides]</p> $\therefore \hat{O}_1 = 2\hat{K}_1$ $\hat{O}_2 = \hat{K}_2 + P \quad [\text{ext } \angle \text{ of } \Delta]$ <p>But $\hat{K}_2 = P$ [\angles opp equal sides]</p> $\therefore \hat{O}_2 = 2\hat{K}_2$ $\therefore \hat{O}_1 + \hat{O}_2 = 2\hat{K}_1 + 2\hat{K}_2$ $= 2(\hat{K}_1 + \hat{K}_2)$ $\therefore \hat{T\hat{O}P} = 2\hat{T\hat{K}P}$ <p>OR</p> | <p>✓ construction</p> <p>✓ S / R</p> <p>✓ S</p> <p>✓ S</p> <p>✓ S</p> <p>(5)</p> |
|------------|--|--|




8.1



| | | |
|-----|---|--|
| 8.1 | <p>Construction: Draw KO</p> $\hat{T} = \hat{K}_1$ <p>[∠ s opp. equal sides]</p> $\therefore \hat{KOT} = 180^\circ - 2\hat{K}_1$ <p>[sum of ∠ s of ΔKOT]</p> $\hat{P} = \hat{K}_2$ <p>[∠ s opp. equal sides]</p> $\therefore \hat{KOP} = 180^\circ - 2\hat{K}_2$ <p>[sum of ∠ s of ΔKOP]</p> $\hat{TOP} = 360^\circ - (\hat{KOT} + \hat{KOP})$ <p>[∠ s around a point]</p> $= 360^\circ - (180^\circ - 2\hat{K}_1 + 180^\circ - 2\hat{K}_2)$ $= 2\hat{K}_1 + 2\hat{K}_2$ $= 2(\hat{K}_1 + \hat{K}_2)$ $\therefore \hat{TOP} = 2\hat{TKP}$ | <p>✓ construction</p> <p>✓ S / R</p> <p>✓ S</p> <p>✓ S</p> <p>✓ S</p> <p>(5)</p> |
|-----|---|--|




8.2



| | | |
|-----|---|----------|
| 8.2 | $\hat{O}_1 = 4x + 100^\circ$ [given] | |
| | $\therefore \hat{A} = 2x + 50^\circ$ [\angle at centre = $2 \times \angle$ at circumference] | ✓ S ✓ R |
| | $x + 34^\circ + 2x + 50^\circ = 180^\circ$ [opp \angle s of cyclic quad] | ✓ S ✓ R |
| | $3x = 96^\circ$ | |
| | $x = 32^\circ$ | ✓ answer |
| | OR | (5) |
| | $\hat{O}_2 = 2x + 68^\circ$ [\angle at centre = $2 \times \angle$ at circumference] | ✓ S ✓ R |
| | $4x + 100^\circ + 2x + 68^\circ = 360^\circ$ [\angle s round a pt] | ✓ S ✓ R |
| | $6x = 192^\circ$ | |
| | $x = 32^\circ$ | ✓ answer |
| | OR | (5) |
| | $\hat{O}_2 = -4x + 260^\circ$ [\angle s round a pt] | ✓ S ✓ R |
| | $2\hat{C} = -4x + 260^\circ$ [\angle at centre = $2 \times \angle$ at circumference] | ✓ S ✓ R |
| | $\hat{C} = -2x + 130^\circ$ | |
| | $x + 34^\circ = -2x + 130^\circ$ | |
| | $3x = 96^\circ$ | |
| | $x = 32^\circ$ | ✓ answer |
| | | (5) |




8.3




| | | |
|-------|---|---------------------------------------|
| 8.3.1 | $\hat{O}MB = 90^\circ$ [∠ in semi circle] | ✓ S ✓ R (2) |
| 8.3.2 | $AB = \sqrt{300} = 10\sqrt{3}$ $\therefore MB = 5\sqrt{3}$ [line from centre \perp to chord] $OB^2 = OM^2 + MB^2$ [Pythagoras] $OB^2 = 5^2 + (5\sqrt{3})^2$ $OB = 10$ units | ✓ S ✓ R ✓ S ✓ answer (4) |
| | | [16] |



QUESTION/VRAAG 9

| | | |
|-----|---|---|
| 9.1 | $\frac{FB}{EB} = \frac{DA}{EA}$ <p>[prop theorem; DC AB] OR [line one side of Δ]</p> $FB = \frac{4p \times 21}{7p}$ $FB = 12 \text{ units}$ | <p>✓ S ✓ R</p> <p>✓ answer</p> <p>(3)</p> |
| 9.2 | <p>In ΔEDF and ΔEAB:</p> <p>\hat{E} is common</p> <p>$\hat{E}DF = \hat{A}$ [corresp \angles; EA CB]</p> <p>$\hat{E}FD = \hat{E}BA$ [corresp \angles; DC AB]</p> <p>$\Delta EDF \parallel \Delta EAB$ [\angle; \angle; \angle]</p> | <p>✓ S</p> <p>✓ S/R</p> <p>✓ S OR R</p> <p>(3)</p> |
| 9.3 | $\frac{DF}{AB} = \frac{ED}{EA}$ <p>[Δs]</p> $DF = \frac{3p \times 14}{7p}$ <p>DF = 6 units</p> <p>FC = 8 units [DC = AB = 14 units; opp sides of ^m]</p> <p>OR</p> <p>$\Delta EDF \parallel \Delta BCF$ [\angle; \angle; \angle]</p> $\frac{ED}{BC} = \frac{DF}{CF}$ <p>[Δs]</p> $\frac{3}{4} = \frac{14 - FC}{FC}$ <p>[BC = AD; opp sides of ^m]</p> $3FC = 56 - 4FC$ <p>FC = 8</p> | <p>✓ S</p> <p>✓ DF = 6</p> <p>✓ FC = 14 – DF</p> <p>(3)</p> <p>✓ $\Delta EDF \parallel \Delta BCF$</p> <p>✓ $\frac{3}{4} = \frac{14 - FC}{FC}$</p> <p>✓ answer</p> <p>(3)</p> |
| | | [9] |

QUESTION/VRAAG 10



| | | |
|------|---|---|
| 10.1 | $\hat{S}_3 = \hat{PQR}$ [ext \angle of cyclic quad] $\hat{R}_3 = \hat{PQR}$ [\angle s opp equal sides] $\therefore \hat{S}_3 = \hat{R}_3$ But $\hat{S}_4 = \hat{R}_3$ [\angle s in the same seg] $\therefore \hat{S}_3 = \hat{S}_4$ | \checkmark S \checkmark R \checkmark S / R \checkmark S \checkmark R (5) |
| 10.2 | $\hat{R}_1 + \hat{R}_2 = \hat{PQR}$ [tan chord theorem] $\hat{S}_4 = \hat{PQR}$ [proved in 10.1] $\therefore \hat{S}_4 = \hat{R}_1 + \hat{R}_2$ SMRC is a cyclic quad [converse ext \angle of cyclic quad] | \checkmark S \checkmark R \checkmark S \checkmark R (4) |
| 10.3 | $\hat{S}_3 = \hat{R}_2 + \hat{P}_2$ [ext \angle of Δ] $\hat{S}_4 = \hat{P}_1 + \hat{A}_2$ [ext \angle of Δ] $\therefore \hat{R}_2 + \hat{P}_2 = \hat{A}_2 + \hat{P}_1$ But $\hat{P}_1 = \hat{R}_2$ [tan chord theorem] $\therefore \hat{P}_2 = \hat{A}_2$ RP is a tangent to the circle [converse tan chord theorem] OR [\angle between line and chord] OR [converse alt seg theorem] | \checkmark S \checkmark R \checkmark S \checkmark S \checkmark R \checkmark R (6) |



| | | |
|---|-------------------------------------|---------|
| In $\triangle MSP$ and $\triangle MPA$ | | |
| \hat{M}_2 is common | | ✓ S |
| $AR = AP$ | [tans from same point] | ✓ S / R |
| $\hat{R}_1 + \hat{R}_2 = \hat{P}_1 + \hat{P}_2$ | [\angle s opp equal sides] | ✓ S |
| $\hat{S}_4 = \hat{R}_1 + \hat{R}_2$ | [proved in 10.2] | |
| $\therefore \hat{S}_4 = \hat{P}_1 + \hat{P}_2$ | | ✓ S |
| $\therefore \hat{P}_2 = \hat{A}_2$ | [sum of \angle s in \triangle] | ✓ S |
| RP is a tangent to the circle | [converse tan chord theorem] | ✓ R |
| | | (6) |
| | | [15] |

TOTAL/TOTAAL: 150