

SA's Leading Past Year

Exam Paper Portal

S T U D Y

You have Downloaded, yet Another Great
Resource to assist you with your Studies ☺

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexamapers.co.za





Province of the
EASTERN CAPE
EDUCATION

Ipondo leMpuma Kapa: Isobe leMfundu
Provincie van die Oos Kaap: Departement van Onderwys
Porafensio Ya Kapa Botjahabelo: Lefapha la Thuto

**NATIONAL
SENIOR CERTIFICATE/
NASIONALE
SENIORSERTIFIKAAT**

GRADE/GRAAD 12

SEPTEMBER 2024

**MATHEMATICS P1/WISKUNDE V1
MARKING GUIDELINE/NASIENRIGLYN**



MARKS/PUNTE: 150

This marking guideline consists of 22 pages./
Hierdie nasienriglyn bestaan uit 22 bladsye.

NOTE/LET WEL:

- If a candidate answers a question TWICE, mark the FIRST attempt ONLY.
Indien 'n kandidaat 'n vraag TWEE keer beantwoord, merk SLEGS die EERSTE poging.
- Consistent accuracy applies in ALL aspects of the marking guideline.
Volgehoue akkuraatheid geld deurgaans in ALLE aspekte van die nasienriglyn.
- If a candidate crossed out an attempt of a question and did not redo the question, mark the crossed-out attempt.
Indien 'n kandidaat 'n poging vir 'n vraag deurgetrek het en nie die vraag weer beantwoord het nie, merk die poging wat deurgetrek is.
- The mark for substitution is awarded for substitution into the correct formula.
- Die punt vir substitusie word toegeken vir substitusie in die korrekte formule.*

QUESTION 1/VRAAG 1

1.1.1	$(2x-4)(x-1)=0$ $x=2 \text{ or/of } x=1$	✓ $x=2$ ✓ $x=1$ (2)
1.1.2	$2x^2 - 3(x+2) = 4$ $2x^2 - 3x - 6 - 4 = 0$ $2x^2 - 3x - 10 = 0$ $x = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(2)(-10)}}{2(2)}$ $x = 3,11 \quad \textbf{OR/OF} \quad x = -1,61$	✓ standard form / standaardvorm ✓ substitution / vervanging ✓ $x = 3,11$ or/of ✓ $x = -1,61$ (4)
1.1.3	$x^2 + 4x - 21 \leq 0$ $(x+7)(x-3) \leq 0$ $c.v's : x \in \{-7; 3\}$  <p style="text-align: center;">OR / OF</p>  <p style="text-align: center;">$\therefore -7 \leq x \leq 3$</p>	✓ factors / faktore ✓✓ answer / antwoord (Accuracy / Akkuraatheid) (3)



1.1.4 $\begin{aligned} -\sqrt{x-1} &= 3-2x \\ 2x-3 &= \sqrt{x-1} \\ (2x-3)^2 &= x-1 \\ 4x^2 - 12x + 9 &= x-1 \\ 4x^2 - 13x + 10 &= 0 \\ (4x-5)(x-2) &= 0 \\ x \neq \frac{5}{4} \text{ or/of } x &= 2 \\ \therefore x &= 2 \end{aligned}$	<ul style="list-style-type: none"> ✓ squaring both sides/ <i>kwadreer beide kante</i> ✓ standard form / <i>standaardvorm</i> ✓ factors / formula <i>faktore / formule</i> ✓ answers with selection <i>antwoorde met keuse</i> <p style="text-align: right;">(4)</p>
---	---





1.3 $f(x) = x^2 + 3x$ $\therefore f(-x) = x^2 - 3x$ $2x = [t(x)]^{\frac{1}{2}}$ $t(x) = 4x^2$ $\therefore t(2k) = 4(2k)^2$ $f(-x) + \frac{t(2k)}{4} = 0$ $x^2 - 3x + \frac{4(2k)^2}{4} = 0$ $x^2 - 3x + 4k^2 = 0$ <p>For equal roots / Vir gelyke wortels, $\Delta = 0$</p> $b^2 - 4ac = 0$ $(-3)^2 - 4(1)(4k^2) = 0$ $9 - 16k^2 = 0$ $(3 - 4k)(3 + 4k) = 0$ $k = \frac{3}{4} \quad \text{or/of} \quad k = -\frac{3}{4}$	$\checkmark \ f(-x) \& t(2k)$ $\checkmark \text{ simplification / vereenvoudiging}$ $\checkmark \text{ subst. in / vervang in: } \Delta = 0$ $\checkmark \text{ method of solving for } k$ $\text{metode vir oplos van } k$ $\checkmark \text{ } k\text{-values / } k\text{-waardes}$	(5) [24]



QUESTION 2/VRAAG 2

2.1		
2.1.1	$\begin{aligned} 2a &= 2 \\ \therefore a &= 1 \\ 3a + b &= 1 \\ b &= 1 - 3(1) \\ \therefore b &= -2 \\ \\ \therefore T_n &= n^2 - 2n - 4 \end{aligned}$	$\begin{aligned} \checkmark a &= 1 \\ \checkmark b &= -2 \\ \checkmark c &= -4 \\ \\ \checkmark T_n &= n^2 - 2n - 4 \end{aligned} \quad (4)$
2.1.2	$\begin{aligned} T_n &= n^2 - 2n - 4 \\ T_{35} &= (35)^2 - 2(35) - 4 \\ &= 1\,151 \end{aligned}$	$\checkmark \text{ answer / antwoord} \quad (1)$
2.1.3	$\begin{aligned} T_n &= 1 + (n-1)(2) \\ T_n &= 2n - 1 \\ T_{n+1} &= 2(n+1) - 1 \\ T_{n+1} &= 2n + 1 \\ T_n \times T_{n+1} &= 1155 \\ (2n-1)(2n+1) &= 1155 \\ 4n^2 - 1 &= 1155 \\ 4n^2 &= 1156 \\ n^2 &= 289 \\ n &= \pm 17 \\ \therefore n &= 17, n \in \mathbb{N} \\ \\ \therefore T_{17} \text{ and/or } T_{18} &\text{ will give a product of 1155} \\ &\text{sal 'n produk van 1 155 gee} \\ &\text{OR/OF} \\ T_n &= 2n - 1 \\ T_{n-1} &= 2n - 3 \\ (2n-1)(2n-3) &= 1155 \\ 4n^2 - 8n - 1152 &= 0 \\ n^2 - 2n - 288 &= 0 \\ (n-18)(n+16) &= 0 \\ n = 18 \text{ or/of } n &\neq -16, n \in \mathbb{N} \\ n = 18 \text{ and/or } n-1 &= 17 \\ \therefore T_{17} \text{ and/or } T_{18} &\text{ will give a product of 1155} \\ &\text{sal 'n produk van 1 155 gee} \end{aligned}$	$\begin{aligned} \checkmark T_n &= 2n - 1 \\ \\ \checkmark (2n-1)(2n+1) &= 1155 \\ \\ \checkmark \text{standard form / standaardvorm} \\ \\ \checkmark n = 17 \text{ and/or } n+1 = 18 \\ \\ \checkmark n = 17, n \in \mathbb{N} \end{aligned} \quad (4)$
		OR/OF
		$\begin{aligned} \checkmark T_n &= 2n - 1 \\ \\ \checkmark (2n-1)(2n-3) &= 1155 \\ \\ \checkmark \text{standard form / standaardvorm} \\ \\ \checkmark n = 18 \text{ and/or } n-1 = 17 \end{aligned}$
		(4)






QUESTION 3/VRAAG 3


3.1.1	$\frac{2}{3}; \frac{2}{9}$	✓ answer / antwoord (1)
3.1.2	$\begin{aligned} S_{\infty} &= \frac{a}{1-r} \\ &= \frac{2}{1-\frac{1}{3}} \\ &= 3 \end{aligned}$	✓ substitution / vervanging ✓ answer / antwoord (2)
3.2	$\sum_{k=3}^m 8(2)^{k-1} = 131\ 040$ $32 + 64 + 128 + \dots$ $r = 2$ $s_n = \frac{a(1 - r^n)}{1 - r}$ $131\ 040 = \frac{32(2^n - 1)}{2 - 1}$ $\therefore 2^n - 1 = 4\ 095$ $2^n = 4\ 096$ $2^n = 2^{12} \quad \text{OR/OF } n = \log_2(4\ 096)$ $\Rightarrow n = 12$ $n = m - 3 + 1$ $12 = m - 2$ $m = 14$	✓ value of a / waarde van a ✓ substitution / vervanging ✓ simplification / vereenvoudiging ✓ value of n / waarde van n ✓ answer / antwoord (5) [8]



QUESTION 4/VRAAG 4

4.1	$x = -5$ $y = -2$	✓ equation of V.A / vergelyking van V.A ✓ equation of H.A / vergelyking van H.A (2)
4.2	$0 = \frac{-1}{x+5} - 2$ $2 = \frac{-1}{x+5}$ $2(x+5) = -1$ $2x = -11$ $x = \frac{-11}{2} = -5,5$ $\therefore \left(-\frac{11}{2}; 0 \right)$	✓ $y = 0$ ✓ answer / antwoord (2)
4.3	$y = \frac{-1}{x+5} - 2$ $y = \frac{-1}{0+5} - 2$ $y = -\frac{11}{5} = -2,2$ $\therefore \left(0; -\frac{11}{5} \right)$	✓ $x = 0$ ✓ answer / antwoord (2)
4.4		✓ both asymptotes / beide asimptote ✓ x-intercept / x-afsnit and/or y-intercept / y-afsnit ✓ shape / vorm (3)
4.5	$y = -(x+5)-2$ $y = -x-7$ OR/OF $y = -x+c$ Subst./Vervang $(-5; -2)$ $-2 = -(-5)+c$ $-7 = c$ $\therefore y = -x-7$	✓ method / metode ✓ answer / antwoord Note: Neem kennis ✓✓ Answer only – Full marks Slegs antwoord - Volpunte (2)

QUESTION 5/VRAAG 5



5.1	$x = \frac{-b}{2a}$ $x = \frac{-(-5)}{2(1)}$ $x = \frac{5}{2} = 2,5$ <p style="text-align: center;">OR / OF</p> $f(x) = x^2 - 5x + 6$ $f'(x) = 2x - 5$ $2x - 5 = 0$ $2x = 5$ $\therefore x = \frac{5}{2} = 2,5$	✓ substitution / vervanging ✓ equation / vergelyking OR / OF ✓ $f'(x) = 0$ ✓ equation / vergelyking (2)
5.2	$f(x) = g(x)$ $x^2 - 5x + 6 = x + 1$ $x^2 - 6x + 5 = 0$ $(x - 5)(x - 1) = 0$ $x = 5 \text{ or/ of } x = 1$ $g(5) = 5 + 1 = 6$ $g(1) = 1 + 1 = 2$ <p style="text-align: center;">B(1; 2) and/en C(5; 6)</p>	✓ $f(x) = g(x)$ ✓ standard form / standaardvorm ✓ x-values / x-waardes ✓ y-values / y-waardes (4)



5.3	$\begin{aligned} h &= x+1-(x^2 - 5x + 6) \\ h(x) &= x+1-x^2 + 5x - 6 \\ &= -x^2 + 6x - 5 \\ x &= \frac{-6}{2(-1)} \\ &= 3 \end{aligned}$ <p style="text-align: center;">OR/OF</p> $\begin{aligned} h'(x) &= -2x + 6 \\ 0 &= -2x + 6 \\ \therefore x &= 3 \end{aligned}$ $h(3) = -(3)^2 + 6(3) - 5 = 4$ <p>\therefore Max. height is 4 units. <i>Maks. hoogte is 4 eenhede.</i></p>	$\checkmark g(x) - f(x)$ $\checkmark h(x)$ $\checkmark x = 3$ \checkmark Max. height / <i>Maks. hoogte</i> (4)
5.4	$\text{Min. of } f : f\left(\frac{5}{2}\right) = \left(\frac{5}{2}\right)^2 - 5\left(\frac{5}{2}\right) + 6 = -\frac{1}{4}$ $\text{Min. of/van } t(x) = -\frac{1}{4} - 2 = -\frac{9}{4}$ $\therefore y \in \left[-\frac{9}{4}; \infty\right); y \in \mathbf{R}$ <p style="text-align: center;">OR/OF</p> $\therefore y \geq -\frac{9}{4}; y \in \mathbf{R}$	$\checkmark \checkmark$ Range of $t(x)$ / <i>Terrein van $t(x)$</i> (2)
5.5	$2 < x < 3$	$\checkmark \checkmark$ $2 < x < 3$ (accuracy / akkuraatheid) (2)
		[14]



QUESTION 6/VRAAG 6

6.1	$f(x) = -\log_c x$ $\frac{1}{2} = -\log_c \left(\frac{1}{2}\right)$ $\sqrt{c} = 2$ $c = 4$ $g(x) = d x^2$ $\frac{1}{2} = d \left(\frac{1}{2}\right)^2$ $d = 2$	✓ subst. of / vervanging van $\left(\frac{1}{2}; \frac{1}{2}\right)$ ✓ value of c / waarde van c ✓ value of d / waarde van d (3)
6.2.1	$g : y = 2x^2$ $g^{-1} : x = 2y^2$ $g^{-1} : y^2 = \frac{1}{2}x$ $g^{-1} : y = \pm \sqrt{\frac{1}{2}x}$ $\therefore y = \sqrt{\frac{1}{2}x}$	✓ swopping x and y omruil van x en y ✓ answer / antwoord (2)
6.2.2	$f(x) = -\log_4 x$ $h(x) = \log_4 x$ $h^{-1}(x) : y = 4^x$	✓ $h(x) = \log_4 x$ ✓ answer / antwoord (2)
6.2.3	$x \in \mathbb{R}$	✓ answer / antwoord (1)
		[8]



QUESTION 7/VRAAG 7

7.1	$A = P(1-i)^n$ $= 180000(1-0,13)^6$ $= R78052,72$	✓ $n = 6$ ✓ substitution / vervanging ✓ answer / antwoord (3)
7.2	$F_v = \frac{x[(1+i)^n - 1]}{i}$ $= \frac{900 \left[\left(1 + \frac{0,08}{12}\right)^{120} - 1 \right]}{\frac{0,08}{12}} + \frac{1300 \left[\left(1 + \frac{0,08}{12}\right)^{60} - 1 \right]}{\frac{0,08}{12}}$ $= 164\,651,4317 + 95\,519,91312$ $= R260171,34$	✓ $n = 120$ and/en $i = \frac{8\%}{12}$ or / of $\frac{8}{1200}$ ✓ $n = 60$ in F ✓ substitution into F_v / vervanging in F_v ✓ answer / antwoord (5)
7.3.1	$OB = P(1+i)^n - \frac{x[(1+i)^n - 1]}{i}$ $OB = 850000 \left(1 + \frac{0,13}{12}\right)^{75} - \frac{9958,39 \left[\left(1 + \frac{0,13}{12}\right)^{75} - 1 \right]}{\frac{0,13}{12}}$ $= R763\,890,54$ <p style="text-align: center;">OR/OF</p> $OB = \frac{x \left[1 - (1+i)^{-n} \right]}{i}$ $OB = \frac{9958,39 \left[1 - \left(1 + \frac{0,13}{12}\right)^{-165} \right]}{\frac{0,13}{12}}$ $= R763\,889,86$	✓ correct substitution into A formula / korrekte vervanging in A formule ✓ correct substitution into Fv formula / korrekte vervanging in Fv formule ✓ answer / antwoord <p style="text-align: center;">OR/OF</p> ✓ $n = 165$ ✓ substitution into the correct formula / vervanging in korrekte formule ✓ answer / antwoord (3)



<p>7.3.2</p> <p>4 Missed instalments / onbetaalde paaiememente</p> <p>OB</p> <p>T₇₅ T₇₆ T₇₇ T₇₈ T₇₉ T₈₀</p> <p>Adjusted instalment / Aangepaste paaiememente</p>	<p>For Outstanding Balance / Vir Uitstaande Balans $= R 763\ 890,54 :$ $A = P(1+i)^n$</p> $A = 763890,54 \left(1 + \frac{0,13}{12}\right)^4$ $A = R 797\ 534,2651$ $P = \frac{x \left[1 - (1+i)^{-n}\right]}{i}$ $797534,2651 = \frac{x \left[1 - \left(1 + \frac{0,13}{12}\right)^{-161}\right]}{\frac{0,13}{12}}$ $x = \frac{797534,2651 \times \frac{0,13}{12}}{\left[1 - \left(1 + \frac{0,13}{12}\right)^{-161}\right]}$ $x = R 10\ 490,96$ <p>∴ Adjusted instalment is R10 490,96</p> <p>OR / OF</p> <p>For Outstanding Balance / Vir Uitstaande Balans $= R 763\ 889,86 :$ $A = P(1+i)^n$</p> $A = 763889,86 \left(1 + \frac{0,13}{12}\right)^4$ $= R 797\ 533,5551$ $797533,56 = \frac{x \left[1 - \left(1 + \frac{0,13}{12}\right)^{-161}\right]}{\frac{0,13}{12}}$ $x = \frac{797533,56 \times \frac{0,13}{12}}{\left[1 - \left(1 + \frac{0,13}{12}\right)^{-161}\right]}$ $x = R 10\ 490,95$	<ul style="list-style-type: none"> ✓ substitution into the correct A formula / vervanging in korrekte A formule ✓ accumulated amount / opgeboude bedrag ✓ substitution into the correct formula / vervanging in korrekte formule ✓ $n = -161$ ✓ answer / antwoord <p>(5)</p> <p>OR / OF</p> <ul style="list-style-type: none"> ✓ substitution into the correct A formula / vervanging in korrekte A formule ✓ accumulated amount / opgeboude bedrag ✓ substitution into the correct formula / vervanging in korrekte formule ✓ $n = -161$ ✓ answer / antwoord <p>(5)</p>
---	--	---

QUESTION 8/VRAAG 8


8.1	$ \begin{aligned} f(x) &= x^2 - 3 \\ f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(x+h)^2 - 3 - (x^2 - 3)}{h} \\ &= \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - 3 - x^2 + 3}{h} \\ &= \lim_{h \rightarrow 0} \frac{2xh + h^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{h(2x + h)}{h} \\ &= \lim_{h \rightarrow 0} 2x + h \\ f'(x) &= 2x \end{aligned} $	<ul style="list-style-type: none"> ✓ substitution into the formula <i>vervanging in die formule</i> ✓ simplification / vereenvoudiging ✓ factorisation / faktorisering ✓ answer / antwoord
8.2.1	$\frac{dy}{dx} = -6x + 7$	<ul style="list-style-type: none"> ✓ $-6x$ ✓ 7
8.2.2	$ \begin{aligned} D_x &\left[\frac{x^3 - 5x^2}{x^3} - \sqrt{x} \right] \\ &= D_x \left[\frac{x^3}{x^3} - \frac{5x^2}{x^3} - x^{\frac{1}{2}} \right] \\ &= D_x \left[1 - 5x^{-1} - x^{\frac{1}{2}} \right] \\ &= 0 + 5x^{-2} - \frac{1}{2}x^{-\frac{1}{2}} \\ &= \frac{5}{x^2} - \frac{1}{2\sqrt{x}} \end{aligned} $	<ul style="list-style-type: none"> ✓ $x^{\frac{1}{2}}$ ✓ $1 - 5x^{-1}$ ✓ 0 & $5x^{-2}$ (zero does not have to be seen) (hoeft nie nul te sien nie) ✓ $-\frac{1}{2}x^{-\frac{1}{2}}$



8.3 $h(x) = -x^3 - 3x^2 + 1$ $g(x) = h'(x)$ $g(x) = -3x^2 - 6x$ <p>Max of $g(x)$ will occur at $g'(x) = 0$ <i>Maks van $g(x)$ sal wees by $g'(x) = 0$</i></p> $g'(x) = -6x - 6 = 0$ $\therefore x = -1$ $g(-1) = -3(-1)^2 - 6(-1)$ $g(-1) = -3 + 6 = 3$ $\therefore \text{largest value} \Rightarrow \text{maximum} = 3$ $\text{grootste waarde} \Rightarrow \text{maksimum} = 3$	$\checkmark g(x) = -3x^2 - 6x$ $\checkmark x = -1$ $\checkmark \text{answer / antwoord}$ (3)	[13]
---	--	-------------






QUESTION 9/VRAAG 9



9.1.3	<p>Point of inflection/Buigpunt/infleksiepunt</p> $x = \frac{x_A + x_B}{2}$ $x = \frac{1+3}{2} = 2$ <p>OR / OF</p> $f'(x) = -9x^2 + 36x - 27$ $f''(x) = -18x + 36$ $f''(x) = 0$ $-18x + 36 = 0$ $18x = 36$ $\therefore x = 2$ $\therefore f(2) = -3(2)^3 + 18(2)^2 - 27(2) = -6$ $\therefore (2; -6)$ <p>Gradient of / Gradiënt van $g(x)$:</p> $f'(2) = -9(2)^2 + 36(2) - 27 = 9$ <p>Gradient of / Gradiënt van $h(x)$:</p> $m_g \times m_h = -1$ $\therefore m_h = -\frac{1}{9}$ $\therefore h(x) = -\frac{1}{9}x$	<p>✓ method / metode</p> <p>✓ $x = 2$</p> <p>✓ $f(2)$</p> <p>✓ $f'(2)$</p> <p>✓ $h(x) = -\frac{1}{9}x$</p>	(5)
9.1.4	<p>$f''(x) > 0$ when $f(x)$ is concave up</p> <p>x-value for point of inflection is 2</p> $\therefore x < 2$ <p>$f''(x) > 0$ wanneer $f(x)$ konkaaf op is</p> <p>x-waarde vir buigpunt/infleksiepunt is 2</p> $\therefore x < 2$	<p>✓✓ answer / antwoord</p>	(2)



<p>9.2</p> <ul style="list-style-type: none"> • $a = 2 > 0$  <ul style="list-style-type: none"> • $(-3; 0) (0; 0) (3; 0)$ (x-intercepts) <i>(x-afsnitte)</i>  <ul style="list-style-type: none"> • x-values of stationary points: <i>x-waardes van stasionäre punte</i> 	<p>✓ shape / vorm</p> <p>✓ intercepts on the graph/ <i>afsnitte op die grafiek</i></p> <p>✓ x- values for stationary points <i>x-waardes vir stasionäre punte</i></p>
	(3)

[17]



QUESTION 10/VRAAG 10

10.1	$S(t) = -3t^2 + 30t$ $S(3) = -3(3)^2 + 30(3)$ = 63 scripts / skrifte	✓ substitution / vervanging ✓ answer / antwoord (2)																				
10.2	$S'(t) = -6t + 30$ For maximum number of scripts, $S'(t) = 0$ / <i>Vir maksimum aantal skrifte, $S'(t) = 0$</i> $-6t + 30 = 0$ $6t = 30$ $t = 5$ (Day 5 / Dag 5)	✓ $S'(t)$ ✓ $S'(t) = 0$ ✓ $t = 5$ (3)																				
10.3	No / Nee <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <th>D1</th><th>D2</th><th>D3</th><th>D4</th><th>D5</th><th>D6</th><th>D7</th><th>D8</th><th>D9</th><th>D10</th></tr> <tr> <td>27</td><td>48</td><td>63</td><td>72</td><td>75</td><td>72</td><td>63</td><td>48</td><td>27</td><td>0</td></tr> </table> Sum/Som = 495 scripts/skrifte	D1	D2	D3	D4	D5	D6	D7	D8	D9	D10	27	48	63	72	75	72	63	48	27	0	✓ No / Nee ✓ explanation / verduideliking (2)
D1	D2	D3	D4	D5	D6	D7	D8	D9	D10													
27	48	63	72	75	72	63	48	27	0													
		[7]																				



QUESTION 11/VRAAG 11

11.1.1	$x = 1 - (0,11 + 0,19 + 0,41)$ $x = 0,29$ $\therefore P(A) = 0,29 + 0,11 = 0,4$	✓ value of x / waarde van x ✓ answer / antwoord (2)
11.1.2	$P(\text{A or/of not/nie B}) = 0,29 + 0,11 + 0,41 = 0,81$	✓✓ answer / antwoord (2)
11.2.1	$a = 4$	✓ answer / antwoord (1)
11.2.2	$\frac{14}{30} = \frac{7}{15}$	✓ answer / antwoord (1)
11.2.3	$P(\text{winning a game}) = \frac{7}{30}$ $P(\text{playing at home}) = \frac{15}{30} = \frac{1}{2}$ $P(\text{winning a game}) \times P(\text{playing at home})$ $= \frac{7}{30} \times \frac{1}{2}$ $= \frac{7}{60} = 0,12$ $P(\text{winning a game and playing at home}) = \frac{3}{30} = 0,10$ $\therefore \text{events are not independent, since}$ $P(\text{winning a game and playing at home}) \neq P(\text{winning a game}) \times P(\text{playing at home})$	✓ $P(\text{winning a game}) \times P(\text{playing at home})$ ✓ $P(\text{winning a game and playing at home})$ ✓ conclusion
	$P(\text{wen 'n wedstryd}) = \frac{7}{30}$ $P(\text{speel tuiswedstryd}) = \frac{15}{30} = \frac{1}{2}$ $P(\text{wen wedstryd}) \times P(\text{speel tuiswedstryd})$ $= \frac{7}{30} \times \frac{1}{2}$ $= \frac{7}{60} = 0,12$ $P(\text{wen wedstryd en tuis wedstryd}) = \frac{3}{30} = 0,10$ $\therefore \text{gebeurtenisse is nie onafhanklik nie, omdat}$ $P(\text{wen wedstryd en speel tuiswedstryd}) \neq P(\text{wen wedstryd}) \times P(\text{speel tuiswedstryd})$	✓ $P(\text{wen wedstryd}) \times P(\text{speel tuiswedstryd})$ ✓ $P(\text{wen wedstryd en speel tuiswedstryd})$ ✓ Gevolgtrekking (3)
		[9]



QUESTION 12/VRAAG 12

12.1	$21 \times 20 \times 10 \times 9 \times 19 \times 18$ $= 12\ 927\ 600$ codes/kodes	✓✓ answer / antwoord (2)																									
12.2	<p>DIGITS / SYFERS:</p> <table style="margin-left: auto; margin-right: auto;"> <tr> <td style="text-align: center;"><u>4</u></td> <td style="text-align: center;"><u>2</u></td> <td style="text-align: center;">or/of</td> <td style="text-align: center;"><u>3</u></td> <td style="text-align: center;"><u>1</u></td> </tr> <tr> <td style="text-align: center;">4</td> <td style="text-align: center;">1</td> <td></td> <td style="text-align: center;">4</td> <td style="text-align: center;">1</td> </tr> <tr> <td style="text-align: center;">6</td> <td style="text-align: center;">2</td> <td></td> <td style="text-align: center;">6</td> <td style="text-align: center;">2</td> </tr> <tr> <td style="text-align: center;">8</td> <td style="text-align: center;">4</td> <td></td> <td style="text-align: center;">8</td> <td style="text-align: center;">4</td> </tr> <tr> <td style="text-align: center;">9</td> <td></td> <td></td> <td style="text-align: center;">9</td> <td></td> </tr> </table> <p>LETTERS BEFORE G / LETTERS VOOR G:</p> <div style="border: 1px solid black; padding: 5px; display: inline-block;"> A; B; C; D; E and/en F </div> <p>Out of 6 letters remove A and E (vowels) = 4 letters <i>Van die 6 letters verwyder A en E (klinkers) = 4 letters</i></p> <p>COMBINED / KOMBINASIE :</p> $n(A) = (4 \times 20 \times 4 \times 2 \times 19 \times 18) + (4 \times 20 \times 3 \times 1 \times 19 \times 18)$ $= 218\ 880 + 82\ 080$ $= 300\ 960$ $P(A) = \frac{n(A)}{n(S)}$ $= \frac{300960}{12927600}$ $= \frac{22}{945}$ $\approx 0,02$	<u>4</u>	<u>2</u>	or/of	<u>3</u>	<u>1</u>	4	1		4	1	6	2		6	2	8	4		8	4	9			9		✓ 4 in $n(A)$ ✓ 4×2 and/en 3×1 in $n(A)$ ✓ dividing by / deel deur 12927600 ✓ answer / antwoord (4) [6]
<u>4</u>	<u>2</u>	or/of	<u>3</u>	<u>1</u>																							
4	1		4	1																							
6	2		6	2																							
8	4		8	4																							
9			9																								

TOTAL/TOTAAL: 150

